

# WEST Search History

DATE: Tuesday, May 20, 2003

<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
side by side			result set
<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=OR</i>			
L11	L10 and l8	24	L11
L10	l5 and ((\$5bacter\$ or lysate).ti. or (\$5bacter\$ or lysate).ab.)	437	L10
L9	L8 and l7	23	L9
L8	L5 and lysate same activat\$5 with (cell or T-cell or b-cell or (t or b or NK or killer) adj3 cell or Nk-t)	196	L8
L7	L6 and (Mycobacter\$3 or Lactobacillus or Bordatella or Cornyebacteria\$3 of Streptococcus or Haemophilus)	321	L7
L6	L5 and (diabetes or autoimmune)	966	L6
L5	(bacter\$ same lysate) and (vaccine or activat\$5 with (cell or T-cell or b-cell or (t or b or NK or killer) adj3 cell or Nk-t))	2962	L5
<i>DB=USPT; PLUR=YES; OP=OR</i>			
L4	(cd25 with cell)	312	L4
L3	pneumococcus with vaccine	56	L3
L2	DPT with vaccine	49	L2
L1	5891435.pn.	1	L1

END OF SEARCH HISTORY

[Home](#)[Help](#)[Subjects](#)[Feedback](#)[Random](#)[Search OMD](#)

---

## diphtheria-tetanus-pertussis vaccine

A vaccine consisting of diphtheria toxoid, tetanus toxoid, and pertussis vaccine. It is usually given to infants three times at two-month intervals, generally at 2, 4, and 6 months of age. The vaccine protects against diphtheria, tetanus, and whooping cough. In most cases the vaccine causes only a temporary fever and discomfort, but in a few cases serious neurological side effects have been observed.

(12 Dec 1998)

---

**Previous:** diphtheria, diphtheria antitoxin, diphtheria antitoxin unit, diphtherial

**Next:** diphtheria toxin, diphtheria toxoid, diphtheric, diphtheritic

---

Published at the Dept. of Medical Oncology, University of Newcastle upon Tyne  
© Copyright 1997-2003 - The CancerWEB Project. All Rights Reserved.

[Home](#)[Help](#)[Subjects](#)[Feedback](#)[Random](#)[Search OMD](#)

---

## pneumococcal vaccine

Vaccine comprised of purified capsular polysaccharide antigen from 23 types of Streptococcus pneumoniae (representing those types responsible for most of the reported pneumococcal diseases in the U.S.).

(05 Mar 2000)

---

**Previous:** pneumococcal pneumonia immunization, pneumococcal polysaccharide, pneumococcal/suppurative keratitis

**Next:** pneumococci, pneumococcal, pneumococcolysis, pneumococcosis

---

Published at the Dept. of Medical Oncology, University of Newcastle upon Tyne  
© Copyright 1997-2003 - The CancerWEB Project. All Rights Reserved.

[Home](#)[Help](#)[Subjects](#)[Feedback](#)[Random](#)[Search OMD](#)

---

## **pertussis vaccine**

A suspension of killed bordetella pertussis organisms, used for immunization against pertussis (whooping cough). It is generally used in a mixture with diphtheria and tetanus toxoids (dtp). There is an acellular pertussis vaccine prepared from the purified antigenic components of bordetella pertussis, which causes fewer adverse reactions than whole-cell vaccine and, like the whole-cell vaccine, is generally used in a mixture with diphtheria and tetanus toxoids.

(12 Dec 1998)

---

**Previous:** pertussis-like syndrome, pertussis syndrome, pertussis toxin, pertussis toxins

**Next:** perula, peruvian, Peruvian bark, Peruvian tarantula, Peruvian wart

---

Published at the Dept. of Medical Oncology, University of Newcastle upon Tyne  
© Copyright 1997-2003 - The CancerWEB Project. All Rights Reserved.